

Specimen Label

WATER SOLUBLE POUCH PREVENTS ADULT MOSQUITO EMERGENCE IN AREAS UP TO 1/8 ACRE (5,445 ft²) FROM A SINGLE APPLICATION POINT

ACTIVE INGREDIENT:

(S)-Methoprene (CAS #65733-16-6):	1.8%
OTHER INGREDIENTS:	98.2%
TOTAL:	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See below for additional precautionary statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters or rinsate.

INTRODUCTION

Aquaprene™ Tossits™ are convenient ready-to-use packets for treating mosquito breeding sites, allowing control of an area up to 1/8 acre (5,445 ft²) in size from a single application point, and larger areas by application of multiple pouches. **Aquaprene Tossits** treatments can be applied prior to flooding (i.e. "pre-hatch") to known mosquito breeding areas, or post-hatch to sites containing mosquito larvae. Treated larvae continue to develop normally to the pupal stage where they die. **Aquaprene Tossits** prevents the emergence of adult mosquitoes from treated sites.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Apply **Aquaprene Tossits** to continuous water mosquito breeding sites up to two feet deep at any time during the mosquito season. **Aquaprene Tossits** is effective when applied pre-hatch, and on 1st, 2nd, 3rd and early 4th instar larval mosquitoes to prevent adult emergence. Under typical environmental conditions, one application will control for up to 7 days. Continue treatments on regular intervals throughout the mosquito season.

Apply **Aquaprene Tossits** prior to flooding (i.e. "pre-hatch") to known mosquito breeding sites. Apply pre-hatch treatments to dry sites up to 21 days prior to flooding to control floodwater mosquitoes.

Apply **Aquaprene Tossits** pre-hatch treatments in the lowest areas of each site to insure water contact

NOTE: **Aquaprene Tossits** are most effective when applied to sites that have continuous (uninterrupted) water surfaces. **Aquaprene Tossits** will not be effective in sites that have isolated pockets of water. The active ingredient S-Methoprene has no effect on mosquitoes that have reached the pupal or adult stage prior to treatment.

The pouches are water sensitive and dissolve when contacted by water. Keep water-soluble pouches sealed in the original foil bag until ready for use. Once outer foil bag is opened, use contents before deterioration from contact with humidity or moisture.

APPLICATION RATES

Select the target mosquito species and corresponding application rate to treat each 1/8 acre area. Apply additional **Aquaprene Tossits** pouches to each 1/8 acre portion in larger sites.

Aedes, Ochlerotatus, Anopheles and Psorophora spp.:

- Apply 1 to 2 **Aquaprene Tossits** pouches to control up to 1/8 acre (5,445 sq. ft.).

Culex, Culiseta, Coquillettidia and Mansonia spp.:

- Apply 2 to 4 **Aquaprene Tossits** pouches to control up to 1/8 acre (5,445 sq. ft.).

Within these ranges, use lower rates when water is shallow (< 2 ft), vegetation and/or organic matter is minimal. Application of **Aquaprene Tossits** to sites subject to water flow or exchange will diminish the product's effectiveness and may require higher application rates within the ranges of the use rates specified.

Specimen Label

APPLICATION SITES

Apply **Aquaprene Tossits** pre-hatch or post-hatch as directed above to temporary and permanent water sites which support mosquito larval development. Examples of typical sites are: freshwater and salt water marshes, cattail marshes, woodland pools and meadows, grassy swales, floodplains, irrigated croplands, pastures and rangeland (without removal of livestock), vineyards, rice fields (including wild rice), fruit and nut orchards, berry fields and bogs, water hyacinth beds, vegetation-choked phosphate pits, wastewater treatment facilities, waste settling ponds, animal waste effluent lagoons, storm water retention basins, retention and detention ponds, drainage areas, roadside ditches, and other natural and manmade depressions where mosquitoes may breed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of the empty outer foil pouch in trash, as long as WSP is unbroken.

WARRANTY AND CONDITIONS OF SALE

Seller makes no warranty, express or implied, concerning the use and handling of this product other than indicated on the label. To the fullest extent permitted by law, buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Always read the label before using this product.

For information, call 1-888-846-7233 or visit our web site: www.aquaprene.com

B2E Biotech LLC, Sag Harbor, New York U.S.A.

EPA Reg No. 75318-10 | U.S. Patent Pending

Aquaprene™ and Tossits™ are trademarks of B2E Biotech LLC

©2006 B2E Biotech LLC | Made in USA

